

Detection of β -lactamase genes and characterization of class 1 integrons in multidrug-resistant *Pseudomonas aeruginosa* strains in Guilan, Iran

Running title: Detection of β -lactamase genes and characterization of class 1 integrons

Tahereh Panahi

Department of Biology, Rasht Branch, Islamic Azad University, Rasht, Iran,
panahi.t22@yahoo.com

ORCID: 0009-0001-5893-9179

Leila Asadpour

Department of Biology, Rasht Branch, Islamic Azad University, Rasht, Iran,
l.asadpour@yahoo.com

ORCID: 0000-0002-8437-0337

Najmeh Ranji

Department of Biology, Rasht Branch, Islamic Azad University, Rasht, Iran,
najmeh.ranji@gmail.com

ORCID: 0000-0001-5233-2427

Corresponding author: Leila Asadpour

Email: l.asadpour@yahoo.com

Tel: +989113383860

Address: Department of Biology, Rasht Branch, Islamic Azad University, Rasht, Iran

Abstract

Background: Infections caused by extended-spectrum β -lactamase (ESBL) producing *Pseudomonas aeruginosa* (*P. aeruginosa*) are a serious concern in hospitals around the world. Many of β -lactamase genes are carried by integrons. This study was conducted to investigate the frequency of β -lactamase genes and characterize class 1 integrons in multidrug-resistant *P. aeruginosa* strains in Guilan, northern Iran.

Methods: A total of 110 *P. aeruginosa* isolates were collected from different hospitals in 2021 and identified using standard microbiological methods. The isolates were studied for their antibacterial susceptibility and ESBL-producing ability by disk diffusion. All ESBL-producing isolates were investigated for the presence of β -lactamase resistant and integron genes by polymerase chain reaction (PCR). Gene cassette screening was done based on sequence analysis of class 1 integrons.

Results: Based on antibiotic susceptibility testing, 40 isolates (37.4%) were ESBL producers. The frequency of β -lactamase genes including *VIM*, *SIM*, *IMP*, *SPM*, and *OXA2* was 10.3%, 1.9%, 20.6%, 14%, and 4%, respectively. *GIM* and *OXA 10* genes were not found in any of the strains. Furthermore, *int1* gene was identified among 37 isolates (34.6%). The sequencing results of *int1* showed 12 different types of gene cassettes among 13 strains. In this assay, *blaOXA-2* was the only bla gene identified in *int1*.

Conclusion: The results of the present study showed that integrons carrying multidrug resistance genes are highly prevalent in *P. aeruginosa* isolates and ESBL genes were also observed in these strains. Therefore, constant monitoring of drug resistance, especially ESBL producers, is critical to disease management in clinical settings.

Key words: *Pseudomonas aeruginosa*, β -lactamase genes, integron, gene cassette

Introduction

Pseudomonas aeruginosa (*P. aeruginosa*) is a Gram-negative bacterium responsible for a wide variety of infections, including urinary tract infection, pneumonia, septicemia, wound infection, meningitis, and many life-threatening infections (1). Various mechanisms such as β -lactamase production, target mutation, overexpression of output pumps, and reduced outer membrane permeability are believed to be involved in the resistance of *P. aeruginosa* to antimicrobial agents (2). Due to the significance of β -lactam antibiotics in hospital use, various forms of β -lactamases, e.g., extended-spectrum β -lactamases (ESBLs), AmpC β -lactamases, and metallo- β -lactamases (MBL), have evolved. ESBLs are the major enzyme in bacteria which are resistant to various β -lactam antibiotics, including penicillins, cephalosporins, and carbapenems. However, they are inhibited by β -lactam inhibitors such as clavulanic acid, sulbactam, and tazobactam (3,4). These enzymes break down the amide bond of the β -lactam ring and inactivate antimicrobial agents. ESBL genes, including *blaSHV*, *blaTEM*, and *blaCTX-M* are encoded by plasmids and can transfer between bacterial species (5,6). These enzymes are highly prevalent in *Enterobacteriaceae* and *P. aeruginosa* strains. Mutations in genes encoding TEM and SHV enzymes lead to the production of ESBLs with an extended substrate catalytic activity allowing for the breakdown of all cephalosporins, penicillins, and aztreonam (7). An increase in the frequency of ESBLs has been reported in recent years, but it varies in different geographical locations. Overall, clinical treatment failure occurs when inappropriate antimicrobial therapy is used to treat infections caused by ESBL-producing *P. aeruginosa* strains. Therefore, enabling laboratory technicians to correctly identify ESBL-producing *P. aeruginosa* isolates would lead to a better selection of antibiotics and improved infection outcomes (8,9). The growing resistance of *P. aeruginosa* to several antibiotics, as a result of excessive antibiotic administration, has resulted in high levels of antibiotic resistance and cross-resistance between antibiotics and the emergence of multidrug-resistant (MDR) forms of *P. aeruginosa* (10). Integrons are genetic elements that can carry multiple antibiotic resistance genes. To date, four classes of integrons have been described in gram-negative bacterial isolates. Three main classes of integrons have a 5' conserved segment, including an *intI* gene encoding an integrase and an *attI* recombination site, but have distinct 3' conserved segments (6). In Class 1 integrons, the 3' conserved segment includes three open reading frames (ORFs)-*qacE* Δ 1, a deletion derivative of the antiseptic resistance gene *qacE*, the *sul1* sulfonamide resistance gene, and ORF5 (of unknown function), or *int* genes, as in Tn402. Gene cassettes are randomly integrated with the region between 3' and 5' of integrons. These cassettes are mobile gene elements that encode one or more antibiotic resistance genes and do not have a promoter sequence in their structure, so the expression of gene cassettes depends on the expression of integrons. Therefore, integrons act both as a gene expression vector and as a natural cloning system (7). More than 80 different gene cassettes of Class 1 integrons have been described and shown to confer resistance to a wide range of antiseptics, disinfectants, and antibiotics such as β -lactams, fluoroquinolones, aminoglycosides, chloramphenicol, trimethoprim, streptothricin, rifampin, erythromycin, fosfomycin, and lincomycin (11). Few studies have been conducted to investigate the prevalence of multidrug-resistant strains of *P. aeruginosa* producing ESBL in Guilan province, northern Iran. A previous study in Guilan Province investigated the antimicrobial resistance, biofilm-forming ability, and virulence potential of *P. aeruginosa* isolated from burn patients and the results of this study showed that 72.2% (65/90) of *P. aeruginosa* isolates were multidrug-resistant (MDR), and 55.5% (50/90) and 35.6% (32/90) were positive for ESBL and MBL production, respectively (12). The present study was conducted to investigate frequency of antibiotic resistance, detection of β -lactamase,

and characterization of gene cassettes carried by integrons in multidrug-resistant *P. aeruginosa* strains in Guilan Province.

Methods

Bacterial Isolates

This descriptive cross-sectional study was performed in 2021. A total of 110 *P. aeruginosa* isolates were collected from different hospitals in Rasht, northern Iran and identified using standard microbiological methods. All clinical samples were inoculated on eosin methylene blue (EMB), MacConkey, and blood agar and incubated at 37°C for 24 hours. *P. aeruginosa* colonies were identified using microbiological methods.

Antibiotic susceptibility testing

Antibiotic susceptibility testing was performed by the Kirby-Bauer disc diffusion method according to the instructions of the Clinical and Laboratory Standards Institute (CLSI, 2020). Antimicrobial discs, including gentamicin (30 µg), amikacin (30 µg), ciprofloxacin (5 µg), co-trimoxazole (25 µg), piperacillin (100 µg), ceftazidime (30 µg), meropenem (10 µg), imipenem (10 µg), colistin (10 µg), azithromycin (30 µg), kanamycin (30 µg), erythromycin (30 µg), ceftriaxone (30 µg), and cefotaxime (30 µg) were used for initial screening of *P. aeruginosa* strains. The ATCC 27853 standard strain of *P. aeruginosa* was used as control. ESBL production was determined by the double disk diffusion method using disks of ceftazidime (30 µg) and cefotaxime (30 µg) alone and in combination with clavulanic acid (10µg) on Muller Hinton agar. A positive test result was defined as a ≥ 5 mm increase in the zone diameter compared to a disk without clavulanic acid. Moreover, an imipenem-EDTA double-disk synergy test was used for MBL enzyme production assay. Enhancements in the diameter of the zone of inhibition in IMP+EDTA in comparison with the IMP-only disks were considered to be MBL producers.

Detection of β -lactamase and integron genes

The DNA of *P. aeruginosa* isolates positive in the DDST were extracted using a DNA extraction kit as instructed by the manufacturer (Sigma-Aldrich, St Louis, MO). Subsequently, the frequency of β -lactamase genes (bla-VIM, IMP, AMPC, SIM, GIM, SPM, OXA10, and OXA2) and *int1* and *Int2* integron genes were determined in strains using PCR with specific primers as shown in Table 1. PCR was carried out in 25 µl PCR volumes containing 10 ng template DNA, 0.5 mM of dNTPs, 10 pm of each primer, and 1 µl of Taq DNA polymerase in 1x PCR buffer. DNA amplification was performed in a Mastercycler Personal thermal cycler (Eppendorf, Germany) based on cycling parameters described in Table 1. PCR products were analyzed in 1% agarose gel containing 25 µg of ethidium bromide in tris-ethylenediaminetetraacetic acid (EDTA) buffer and the gel was photographed under an ultraviolet illuminator using a gel documentation system and confirmed by sequencing.

Detection of gene cassettes in *int1*

Gene cassettes within class 1 integrons were amplified using 5'-CS and 3'-CS primer pairs (Table 1). The PCR products were then sequenced and identified using the basic local alignment search tool (BLAST) for the gene cassette screening strategy. For PCR products longer than 700 bp, internal primers were designed based on the obtained sequences and the internal parts of the integrons were amplified and subsequently sequenced.

Table 1. Nucleotide sequences of primers used in this study

Gene	Primer sequence (5'-3')	Annealing temperature (°C)	Ref.
<i>VIM</i>	F: GTGTTTGGTCGCATATCGC R: CGCAGCACCAGGATAGAAG	53	13
<i>IMP</i>	F: GGAATAGAGTGGCTTAATTC R: GCCAAGCTTCTATATTTGCG	55	13
<i>SIM</i>	F: TACAAGGGATTTCGGCATCC R: TAATGGCCTGTTCCCATG	52	14
<i>GIM</i>	F: TCGACACACCTTGGTCTG R: AACTTCCAACCTTTGCCAT	52	14
<i>SPM</i>	F: CCTACAATCTAACGGCGACC R: TCGCCGTGTCCAGGTATAAC	53	14
<i>OXA2</i>	F: ATGGCAATCCGAATCTTCGC R: TTATCGCGCAGCGTCCGAGT	55	13
<i>Int1</i>	F: CAGTGGACATAAGCCTGTTC R: CCGAGGCATAGACTGTA	55	9
<i>In-(5'CS)</i>	F: GGCATCCAAGCAGCAAGC R: AAGCAGACTTGACCTGAT	52	9

Results

Antibiotic resistance profile of *P. aeruginosa* strains

A total of 110 *P. aeruginosa* isolates were recovered during the study period. Based on antibiotic susceptibility testing, the highest phenotypic resistance was against ceftazidime and ceftazidime. Meanwhile, colistin and piperacillin were the most effective antibiotics. More than 85% of the isolates (n = 90) were multidrug-resistant (MDR) phenotype (resistant to three classes of antibiotics) and 37% were extensively drug-resistant (XDR) phenotype (resistant to three classes of antibiotics plus imipenem). Seven isolates were resistant to all evaluated antibiotics. Moreover, according to DDST, 40 isolates (37.4%) were ESBL-positive and 28 isolates (25.45%) were MBL-positive. The results of antibacterial resistance are presented in Table 2.

Table 2. Antimicrobial resistance properties in *Pseudomonas aeruginosa* strains

Antibiotic	Resistant isolates
Amikacin	56 (52.33%)
Gentamycin	49 (45.7%)
Ceftazidime	62 (58%)
Cefotaxime	36 (33.6%)
Ceftriaxone	41 (38.3%)
Cefoxitin	87 (81.3%)
Imipenem	45 (42.0%)
Meropenem	36 (33.6%)
Piperacillin	16 (15.0%)
Co-Trimoxazole	25 (23.3%)
Erythromycin	40 (37.4%)
Azithromycin	29 (27.1%)
Enrofloxacin	49 (45.8%)
Ciprofloxacin	55 (51.4%)

Frequency of β -lactamase and class I integron genes

β -lactamase genes, including *bla-VIM*, *IMP*, *SIM*, *GIM*, *SPM*, *OXA10*, and *OXA2* were detected in ESBL- and MBL-positive strains. Based on the results of PCR, the frequency of *VIM*, *SIM*, *IMP*, *SPM*, and *OXA2* were 10.3%, 14%, 20.6%, 1.9%, and 4%, respectively. *OXA10*, *SPM*, and *GIM* genes were not detected in any of the isolates. Furthermore, *int1* gene was identified in 37 isolates (34.6%). *Int2* gene was not detected in any of the isolates.

Characterization of *int1* gene

Primers related to conserved regions of *int1* gene were used for the detection of *int1* polymorphism in *P. aeruginosa* strains. PCR results showed bands with lengths between 700 and 3000 bp (Figure 1).




Figure 1. Size polymorphism in class 1 integron gene amplicons. Lanes M; 100bp molecular marker, Lanes 1-10; Size polymorphism in class 1 integron gene amplicons in tested isolates.

Identification of *int1* gene cassettes

Among 13 selected isolates, 12 different types of gene cassettes in class 1 integron and 18 different genes were identified in sequence analysis of *int1* gene. The most frequent genes were *dfrA1* (n = 5), *aadA2* (n = 4), and *aacA4* (n = 4). The class D β -lactamase gene, *blaOXA-2*, was the only *bla* gene identified in examined isolates. The number of gene cassettes in tested isolates varied between 1-4. The type of gene cassettes identified in class 1 integrons among *P. aeruginosa* isolates is shown in Table 3.

Table 3. Class 1 integron associated gene cassettes identified in *Pseudomonas aeruginosa* strains

Type of gene cassette	Number of isolates
<i>dfrA17</i>	1
<i>aadA1</i>	1
<i>aadB</i>	2
<i>aacA4</i>	1
<i>dfrA1</i>	1
<i>aacA4, dfrA1</i>	1
<i>aadB, dfrA1</i>	1
<i>dfrA32, aadA2, ereA</i>	1
<i>aadA2, aadA, ereA</i>	1
<i>aadA, ereA, aadA2</i>	1
<i>aacA4, dfrA1, aadA2</i>	1
<i>aacA8, blaOXA-2, aacA7, aadA6</i>	1

Discussion

P. aeruginosa, one of the most common nosocomial infectious agents, has led to the death of a large number of hospitalized patients (15). ESBL antibiotics are commonly used for the treatment of diseases caused by these Gram-negative bacteria. However, some bacteria are resistant to β -lactam antibiotics due to ESBL production (16). In the current study, the rate of MDR *P. aeruginosa* was 85%. *P. aeruginosa*, especially its MDR strains, is very common in Iranian hospitals. Similar resistance rates were reported in other researches in Iran. For instance, Nikokar et al. documented a rate of 45.3% multi-drug resistance among burn patients in Guilan in 2012 (17). In 2017, Fazeli et al. evaluated patients admitted to a teaching hospital and reported a rate of 73% multi-drug resistance (18). In 2015, Ghanbarzadeh et al. (2015) conducted a single-center study on burn patients in Tehran and estimated the multi-drug resistance rate at 93.1% (19). In our study, the highest and lowest rates of resistance belonged to cefoxitin (81.3%) and piperacillin (15%), respectively. Rates of resistance to ceftazidime and ceftriaxone were 62% and 56%, respectively. Moreover, our results suggested high resistance to imipenem at 42.05% and the rate of resistance to cefotaxime was 33.6%. The increase in resistance to cephalosporins and imipenem is concerning. This might be due to selective pressure to these antibiotics and it is highly important to monitor the prescription of these antibiotics. In accordance with present results, Ameen et al. investigated the antibiotic resistance profile in *Pseudomonas aeruginosa* strains and reported that among 230 strains of *P. aeruginosa*, 49.5% were resistant to imipenem (20).

In the present study, 37% of test isolates showed XDR phenotype, 37.4% of isolates were ESBL-positive and 25.45% were MBL-positive based on DDST. The emergence of ESBL-producing *P. aeruginosa* is increasingly reported as a major cause of health-care associated infections (3). Overuse of antibiotics can lead to more resistant strains resulting in gene and horizontal diffusion. High frequencies of ESBL production in *P. aeruginosa* isolates from burn patients in Iran were previously reported by Rafiee et al. (39.2% ESBL and 37.3% MBL production among 51 *P. aeruginosa* isolates) (21). This study demonstrated that the most prevalent genotype for MBL production was *blaIMP*, which was detected in 20.6% of isolates, followed by *SIM* (14%), *VIM* (10.3%), *SPM* (1.9%), and *OXA2* (4%). *OXA 10* and *GIM* genes were not observed in any of the strains. The *blaVIM* and *blaIMP* genes were common in MDR isolates of *P. aeruginosa* in previous researches in Iran (22,23). On the other hand, a similar study in South Africa reported low frequency of *blaIMP*, while *blaVIM* was not detected among *P. aeruginosa* isolates (3). Moreover,

in accordance with our results, *blaIMP* was the most frequently detected metallo- β -lactamase gene in a study conducted by Haghi et al. (9). The prevalence of class 1 integrons in this study was 34.6%, which is similar to the findings of Khosravi et al. in Ahvaz (24). The gene cassettes identified in 13 selected isolates included 1-4 different genes in class 1 integron. A class 1 integron carrying the carbapenem resistance gene (*aacA8-blaOxa-2-aadA7-aadA6*) was also identified in meropenem-resistant *P. aeruginosa*. A high frequency of aminoglycoside resistance gene cassettes including *aadA* family, which induce streptomycin and spectinomycin resistance, was identified in this assay. Gene cassettes *dfrA1*, *aad8*, *dfrA17*, *dfrA32*, and *aacA8* were identified for the first time in this study. High frequency of aminoglycoside resistance gene cassettes in class 1 integrons has been reported in several studies. In a previous report, different aminoglycoside resistance genes, including *aadA1*, *aadA2*, *aadA5*, *aadA6*, *aadB*, *accA4*, and *aac(6)-IIa*, and three *bla* genes *blaOXA10*, *blaP1*, and *blaCARB-8* were identified in integrons structure (25).

In a study conducted by Ahmadian et al. in north of Iran, the resistance genes identified in class I integrons included: *aadA6-orfD* (35.71%), *aacA4-blaOXA-10* (21.42%), *aadB-aacA4-blaOXA-10* (19.04%), *blaOXA-10-aacA4-VIM1* (11.9%), *aacA4-catB10* (7.14%), *aacA5-aadA1-cmlA5* (7.14%), *blaOXA31-aadA2* (4.76%), and *aac(3)-Ic-aacA5-cmlA5* (4.76%), *blaOXA-10-aacA4-VIM1* cassette array and aminoglycoside resistance genes were predominant (26). The highest frequency belonged to *aadA2*, which is similar to our findings. In another study from Iran, analysis of integron cassettes sequence indicated the presence of *dfrA17*, *dfrA7*, *aadB*, *aadA1*, and *dhfr1-sat2* resistance gene cassettes among the isolates (9).

The predominant cassette arrangement described by Hsiao et al. among 11 different types of gene cassette arrays in *P. aeruginosa* isolates in southern Taiwan included *aac(6')-II-catB2-aadA2*, *aac(6')-II-aadA2*, *aac(6')-II-catB2*, *aacA4-aadA15*, *aacC1-orfA-orfB-aadA1*, *cmlA-aadA1*, *catB3-blaOxa-10-aadA15*, *aacA4-catB8-aadA1*, *aadB-orfF1-aadA11*, *dfrB1*, and *dfrB4a-aacA4-aacA4-aadA1* (27). In 2018, Chowdhury et al. examined *P. aeruginosa* isolates with widespread antibiotic resistance in Australia and identified two genomic islands that carried transposons Tn6161 and Tn6163, respectively. A genomic island *GII* gene cassette *blaGES-5-aacA4-gcuE15-aphA15* was also found in class 1 integron, which confirms resistance to carbapenems and aminoglycosides (28). The presence of *aacA4* gene cassette in our study was similar to the findings of Chowdhury et al. (28). In addition, high prevalence of *aadA* gene, which shows resistance to streptomycin/spectinomycin, was observed in a study conducted by Bamidele et al. Gene cassettes identification showed the presence of *aadA6-orfD* and *aadA13*, respectively (29).

In the present study, the investigated class 1 integrons included 12 gene cassettes: *aadA2*, *aadA6*, *era*, *aacA4*, *dfrA1*, *aad8*, *aacA8*, *blaOxa2*, *aacA7*, *dfrA17*, *dfrA32*, and *aadB*. Gene cassettes *dfrA1*, *aad8*, *dfrA17*, *dfrA32*, and *aacA8* were first identified in this study. In a study in China, many genes, including *aadA1*, *aadA2*, *aadA5*, *aadA6*, *aadB*, *accA4*, and *aac(6')-IIa* in integron structures were associated with aminoglycoside resistance. Resistance genes that are often identified (*aadA* family) are adenylyl transferase aminoglycoside genes that cause resistance to streptomycin and spectinomycin. In addition, three genes, i.e., *blaOXA10*, *blaP1*, and *blaCARB-8*, were identified inside integrons (25). Gene cassettes *aadB*, *accA4*, *aadA1*, and *aadA2* in the mentioned study are similar to those found in our study.

Conclusion

The high prevalence of class 1 integrons and resistance genes is a serious concern. It might jeopardize the control and treatment of Pseudo-formal infections, increase the length of hospital stay, and raise mortality rates. Appropriate measures must hence be taken to ensure the proper use

of antibiotics in clinical and agricultural settings. In general, the results of the present study showed that class 1 integrons are widely distributed among *P. aeruginosa* strains isolated from Guilan hospitals. They can be considered as carrier of gene cassettes and can play an important role in the acquisition of MDR in this bacterium.

Acknowledgments

Support by Islamic Azad University, Rasht Branch is gratefully acknowledged.

Conflict of interest

All contributing authors declare no conflicts of interest.

Ethics approval

Since we did not use any animal models and patients and just used isolates that were previously obtained from clinical samples in laboratories, we didn't have any ethical approve for this study but we confirm that the study complies with all regulations.

Funding source

None.

References

- 1- Qin S, Xiao W, Zhou C, Pu Q, Deng X, Lan L, Liang H, Song X, Wu M. *Pseudomonas aeruginosa*: pathogenesis, virulence factors, antibiotic resistance, interaction with host, technology advances and emerging therapeutics. *Signal Transduct Target Ther*. 2022;7(1):199.
- 2- Pang Z, Raudonis R, Glick BR, Lin TJ, Cheng Z. Antibiotic resistance in *Pseudomonas aeruginosa*: mechanisms and alternative therapeutic strategies. *Biotechnology advances*. 2019 Jan 1;37(1):177-92.
- 3- Hosu MC, Vasaikar SD, Okuthe GE, Apalata T. Detection of extended spectrum beta-lactamase genes in *Pseudomonas aeruginosa* isolated from patients in rural Eastern Cape Province, South Africa. *Scientific reports*. 2021 Mar 29;11(1):1-8.
- 4- Başkan C, Sırıken B, Tüfekci EF, Kılınç Ç, Ertürk Ö, Erol İ. Presence of quorum sensing system, virulence genes, biofilm formation and relationship among them and class 1 integron in carbapenem-resistant clinical *Pseudomonas aeruginosa* isolates. *Arch Microbiol*. 2022; 204(8):464.
- 5- Büchler AC, Wüthrich D, Wicki Jauslin M, Egli A, Widmer AF. Nosocomial transmission of a blaVIM-2 carbapenemase integron between isolates of two different *Pseudomonas* species. *Infect Control Hosp Epidemiol*. 2022;43(2):245-247
- 6- El Sayed Zaki M, Mostafa Mahmoud N, Anies Rizk M. Molecular Study of Integrase Gene I and Integrase Gene II in Clinical Isolates of *Pseudomonas aeruginosa*. *Infect Disord Drug Targets*. 2022 Apr 8. doi: 10.2174/1871526522666220408111454.
- 7- Abd-Elmonsef MME, Maxwell SY. Class 1, 2 and 3 integrons in clinical *Pseudomonas aeruginosa* isolated from Tanta University Hospitals, Egypt. *J Chemother*. 2022;34(4):241-246.
- 8- Uddin F, Sohail M, Shaikh QH, Hussain MT, Roulston K, McHugh TD. Verona integron-encoded metallo-Beta-lactamase (VIM) and Vietnam extended-spectrum Beta-lactamase (VEB)

producing *Pseudomonas balearica* from a clinical specimen. J Pak Med Assoc. 2022 Apr;72(4):761-763.

9- Haghi F, Keramati N, Hemmati F, et al. Distribution of Integrins and gene cassettes among metallo- β -lactamase producing *Pseudomonas aeruginosa* clinical isolates. Infect Epidemiol Med. 2017; 3(2): 36-40.

10- Brovedan MA, Marchiaro PM, Díaz MS, Faccone D, Corso A, Pasteran F, Viale AM, Limansky AS. *Pseudomonas putida* group species as reservoirs of mobilizable Tn402-like class 1 integrons carrying blaVIM-2 metallo- β -lactamase genes. Infect Genet Evol. 2021; 96:105131.

11- Aryanezhad M, SHakibaie MR, Karmostaji A, et al. Prevalence of Class 1, 2 and 3 Integrins and Biofilm Formation in *Pseudomonas aeruginosa* and *Acinetobacter baumannii* among ICU and non- ICU Patients, Infect Epidemiol Med. 2016 Autumn; volume 2, issue 4:1-7.

12- Asadpour L. Antimicrobial resistance, biofilm-forming ability and virulence potential of *Pseudomonas aeruginosa* isolated from burn patients in northern Iran. J Glob Antimicrob Resist. 2018; 1(13):214-220.

13- Garza-Ramos U, Morfin-Otero R, Sader HS, et al. Rodriguez-Noriega, A. Sanchez, B. Carrillo, S. Esparza-Ahumada, and J. Silva-Sanchez Metallo- β -Lactamase Gene blaIMP-15 in a Class 1 Integron, In95, from *Pseudomonas aeruginosa* Clinical Isolates from a Hospital in Mexico, Antimicrob Agents Chemother. 2008 Aug; 52(8): 2943–2946.

14- Azimi A, Peymani A and Kianoush pour P. Phenotypic and molecular detection of metallo- β -lactamase-producing *Pseudomonas aeruginosa* isolates from patients with burns in Tehran, Iran, Rev Soc Bras Med Trop 51(5):610-615, Sep-Oct, 2018.

15- Emami S, Nikokar I, Ghasemi Y, et al. Antibiotic Resistance Pattern and Distribution of *pslA* Gene Among Biofilm Producing *Pseudomonas aeruginosa* Isolated From Waste Water of a Burn Center. Jundishapur J Microbiol 2015; 8: e23669.

16- Mirsalehian A, Feizabadi M M., Nakhjavani F A, et al. Detection of VEB-1, OXA-10 and PER-1 genotypes in extended-spectrum-lactamase-producing *Pseudomonas aeruginosa* strains isolated from burn patients, February 2010 Burns: journal of the International Society for Burn Injuries 36(1):70-74.

17- Nikokar I, Tishayar A, Flakiyan Z, Alijani K, Rehana-banisaeed S, Hossinpour M, et al. Antibiotic resistance and frequency of class 1 integrons among *Pseudomonas aeruginosa*, isolated from burn patients in Guilan, Iran. Iran J Microbiol 2013; 5(1): 36-41.

18- Fazeli H, Fatahi Bafghi M, Faghri M, Akbari R. Molecular study of PER and VEB genes in multidrug resistant *Pseudomonas aeruginosa* isolated from clinical specimens in Isfahan/Iran and their antibiotic resistance patterns. J Kerman Univ Med Sci 2012; 19(4): 345-53.

19- Ghanbarzadeh Corehtash Z, Ahmad Khorshidi FF, Akbari H, Aznavah AM. Biofilm formation and virulence factors among *Pseudomonas aeruginosa* isolated from burn patients. Jundishapur J Microbiol. 2015;8:e22345.

20- Ameen N, Memon Z, Shaheen S, Fatima G, Ahmed F. Imipenem Resistant *Pseudomonas aeruginosa*: The fall of the final quarterback. Pak J Med Sci. 2015;31(3):561-5. doi: 10.12669/pjms.313.7372. PMID: 26150844; PMCID: PMC4485271.

21- Rafiee R, Eftekhari F, Tabatabaei SA, Minaee Tehrani D. Prevalence of extended-spectrum and metallo- β -lactamase production in AmpC β -lactamase producing *Pseudomonas aeruginosa* isolates from burns. Jundishapur J Microbiol. 2014; 7 (9):e16436.

22- Anvarinejad M, Japoni A, Rafeatpour N, et al. Burn patients infected with Metallo-Beta-Lactamase-producing *Pseudomonas aeruginosa*: multidrug-resistant strains. Arch Trauma Res 2014; 3: e18182.

- 23-Fallah F, Shams Borhan R, Gholinejad Z, Zahirnia Z, Adabiyani S, Sattarzadeh Tabrizi M et al . Detection of blaIMP and blaVIM Metallo-Beta-Lactamases Genes in *Pseudomonas aeruginosa* Strains Isolated from Wound of Burnt Patients in Tehran Shahid Motahari Hospital during 2011, Iran. Qom Univ Med Sci J. 2013; 7 (5) :21-27
- 24- Khosravi AD, Motahar M, Abbasi MontazeriE (2017) The frequency of class1 and 2 integrons in *Pseudomonas aeruginosa* strains isolated from burn patients in a burn center of Ahvaz, Iran. PLoS ONE 12(8): e0183061.
- 25-Gu B, Tong M, Zhao W, Liu G, Ning M, Pan S, et al. Prevalence and characterization of class 1 integrons among *Pseudomonas aeruginosa* and *Acinetobacter baumannii* isolates from patients in Nanjing, China. J Clin Microbiol 2007; 45(1): 241- 3.
- 26-Ahmadian L, Haghshenas M, Mirzaei B, et al. Distribution and Molecular Characterization of Resistance Gene Cassettes Containing Class 1 Integrons in Multi-Drug Resistant (MDR) Clinical Isolates of *Pseudomonas aeruginosa*, Mazandaran,Iran, Infection and Drug Resistance,2020; 13: 2773–2781.
- 27-Hsiao K, Lee M, Peng C. Detection and characterization of class 1integron-associated gene cassettes from *Pseudomonas aeruginosa* isolates in southern Taiwan. Biomarkers and Genomic Medicine (2014) 6, 74-78.
- 28-Chowdhury V, Gunjal S, Mehta M. Antibiotic resistance patterns of *Pseudomonas aeruginosa* in a tertiary care hospital in Central India. International Journal of Medical Science and Public Health. Int J Med Sci Public Health. 2013; 2(2). 386-389.
- 29- Bamidele T, Odumosu, Bolanle A, et al. Analysis of integrons and associated gene cassettes in clinical isolates of multidrug resistant *Pseudomonas aeruginosa* from Southwest Nigeria. Odumosu et al. Annals of Clinical Microbiology and Antimicrobials 2013.