



Does Physical Activity Level Affect Homocysteine and Obesity Variables in Women with Cardiovascular Disease?

Toktam Arbabian

(PhD Candidate) Department of Physical Education and Sport Sciences, Aliabad Katoul Branch, Islamic Azad University, Aliabad Katoul, Iran

Reza Rezaeshirazi

(PhD) Department of Physical Education and Sport Sciences, Aliabad Katoul Branch, Islamic Azad University, Aliabad Katoul, Iran

Amin Farzaneh Hesari

(PhD) Department of Physical Education and Sport Sciences, Sari Branch, Islamic Azad University, Sari, Iran

Habib Asgharpoor

(PhD) Department of Physical Education and Sport Sciences, Aliabad Katoul Branch, Islamic Azad University, Aliabad Katoul, Iran

Corresponding author: Reza Rezaeshirazi

Tel: +9891 12767500

Email: dr.rezaee@aliabadiu.ac.ir

Address: Department of Physical Education and Sport Sciences, Aliabad Katoul Branch, Islamic Azad University, Aliabad Katoul, Iran

Received: 2020/08/04

Revised: 2020/09/01

Accepted: 2020/09/29



© The author(s)

ABSTRACT

Background and objectives: Physical activity has long been considered as the cornerstone of interventions for reducing the burden of cardiovascular diseases (CVDs). Homocysteine and obesity indices are important indicators of CVD risk. The present research aimed to evaluate homocysteine and obesity variables in female CVD patients and healthy counterparts with different physical activity levels.

Methods: This study was carried out on 85 CVD patients (40-55 yr, n=41, 60-75 yr, n=44) and 80 healthy women (40-50 yr, n=40, 60-75 yr, n=40). Subjects were divided into three groups based on their physical activity level: sedentary, moderately active and active. Serum homocysteine, body roundness index (BRI), a body shape index (ABSI), body mass index (BMI) and waist circumference (WC) were measured. Data were analyzed using the independent t-test and one-way ANOVA.

Results: Physically active middle-aged CVD patients had significantly lower BRI ($p=0.022$) and homocysteine ($p=0.008$) levels compared with the sedentary counterparts. In addition, physically-active old CVD patients had significantly lower BRI ($p=0.041$), ABSI ($p=0.011$) and homocysteine ($p=0.001$) compared with the sedentary counterparts. Physically-active healthy middle-aged individuals had significantly lower BRI ($p=0.013$) and BMI ($p=0.008$) levels compared with sedentary counterparts. Active elderly subjects also had significantly lower BRI ($p=0.001$) and WC ($p=0.008$) compared with the sedentary counterparts.

Conclusion: Physical activity can reduce the risk of CVD and improve health status of women by reducing homocysteine, BRI and ABSI.

Keywords: [Exercise](#), [Cardiovascular Diseases](#), [Homocysteine](#), [Obesity](#).

INTRODUCTION

Cardiovascular diseases (CVDs) are serious disorders affecting the heart and blood vessels that are also the main cause of mortality worldwide (1). According to the World Health Organization (WHO), lack of physical activity is a major risk factor for the development of CVD. This group of disorders is known as the fourth leading cause of mortality and morbidity in the world. Regular physical activity in adults decreases risk of hypertension, coronary artery diseases (CAD) and stroke (2). Unlike some CVD risk factors that have a genetic basis and are not modifiable (e.g. hypertension and hyperlipidemia), low physical activity could be resolved by proper planning as well as change in attitude and behavior (2).

It is established that adults characterized by excessive adipose tissue and ectopic fat stores have a higher risk of CVDs (3). Evidence indicates an association between obesity and increased risk of hypertension, type-2 diabetes mellitus and dyslipidemia, which are major risk factors for CVD (4). Body mass index (BMI) is generally recognized as an anthropometric index of obesity that might also predict the risk of CVD (5). Nevertheless, the neglect of body fat distribution is considered as a typical limitation of BMI. Waist circumference (WC) is recommended as an effective index for measuring visceral fat distribution (6). However, none of these indices alone is able to distinguish subcutaneous fat from visceral fat, which is more strongly associated with metabolic abnormalities. Some studies still suggest that these indices provide limited information on fat distribution (7).

Recently, a body shape index (ABSI) was proposed by Krakauer and Krakauer (8) as a new body index that is more significantly associated with abdominal adipose tissue and premature death than WC and BMI (9,10). However, some studies reported that ABSI seemed to be a weaker index for identifying risk of hypertension, diabetes, CVD and metabolic syndrome compared with BMI (11,12). Body roundness index (BRI) is another index recently proposed by Thomas et al. and based on WC and height. This index is a good predictor of body fat percentage and visceral adipose tissue compared with the traditional indices such as BMI, WC or hip circumference (13). Several studies have

shown that BRI could be used as an indicator of adipose distribution and the presence of eccentric left ventricular hypertrophy, hyperuricemia, CVD and diabetes (11,14).

Homocysteine (Hcy) is a sulfhydryl-containing amino acid produced via methionine demethylation, which is essential for intravascular metabolism (15). Hyperhomocysteinemia has been considered as a risk factor for systemic atherosclerosis, CVD and stroke (16). However, results of studies on the relationship of physical activity with Hcy concentration are contradictory (17-19). Given the importance of physical activity and its role in the prevention of chronic diseases, the aim of the present study was to evaluate Hcy, BRI and ABSI levels in CVD women with different physical activity levels.

MATERIALS AND METHODS

The study was carried out on 85 patients with symptoms of CVD (middle-aged: 40-55 yr, n=41 and elderly: 60-75 yr, n=44) who had been referred to the heart unit of hospitals in Mashhad (Iran) in 2019. The study received approval from the ethics committee of Islamic Azad University, Aliabad Katool branch, Iran (ethics code: IR.IAU.AK.REC.1398.013). A written informed consent was obtained from all subjects before participating in the study. Participants were categorized into two groups: CVD (myocardial infarction, stroke and heart failure) and CVD risk factors (hypertension and hypercholesterolemia).

For middle-aged and elderly groups, an age-matched control group comprising of healthy individuals was also included. Individuals with history of smoking and kidney disorders were excluded. Patients and healthy subjects were divided into three groups based on their physical activity level: sedentary, moderately active and active. A questionnaire was also used to record the participants' dietary habits over the past year.

Physical activity assessment

The International Physical Activity Questionnaire (IPAQ) was used for determining the physical activity. Weekly physical activity was computed by multiplying time (minutes of given activity in the reported week) by intensity (MET) corresponding to that activity. The total weekly activity was obtained by calculating the sum of the activities and expressed in MET·min/week.

Based on the results, the subjects were classified as sedentary (score of less than 600 MET·min/week), moderately active (score of at least 600 but less than 1600 MET·min/week) and active (score of more the 1600 MET·min/week) (20).

Obesity variables

Weight, height, WC and BMI were measured according to standard procedures. The ABSI was calculated using the following formula

$$(8): \text{ABSI} = \left(\frac{WC}{\text{BMI}^{2/3} \times \text{height}^{1/2}} \right)$$

The BRI was based on height (m) and waist circumference (m). First, the eccentricity of the body was determined using the formula below. Eccentricity (non-dimensional value) quantifies the degree of circularity of an ellipse, and it ranges between zero (perfect circle) to one (a vertical line) (14).

$$\varepsilon = \sqrt{1 - \left(\frac{(WC/(2\pi))^2}{(0.5 \times \text{height})^2} \right)}$$

Subsequently, BRI was calculated using the formula below. As described by Thomas et al., values closer to 1 are related to leaner individuals, whereas larger values are associated with rounder individuals (13).

$$\text{BRI} = 364.2 - (365.5 \times \varepsilon)$$

Blood sample collection

Blood samples (5 ml) were taken from subjects' forearms in a comfortable sitting position after at least eight hours of overnight fasting. The samples were centrifuged at 2000 rpm for 10 min to separate serum. Serum samples were then stored at 70 °C until use. The level of serum Hcy was measured using a commercial ELISA kit (Axis-shield diagnost, Germany) according to the manufacturer's

instructions.

Statistical analysis

The Shapiro-Wilk was used to assess normal distribution of data. The independent t-test and one-way analysis of variance (ANOVA) with Tukey post-hoc test were used for analysis of data. All statistical analyses were performed in SPSS24 and at significance of ≤ 0.05 .

RESULTS

The basic characteristics and mean of variables in different groups for middle-aged subjects are shown in table 1.

Compared with healthy subjects, CVD patients had higher WC, BMI, Hcy, BRI and ABSI levels. Physically-active CVD patients had significantly lower BRI ($p=0.022$) and Hcy ($p=0.008$) levels compared with sedentary subjects. There was no significant difference in ABSI, BMI and WC between CVD patients with different activity levels.

Active healthy subjects had significantly lower BRI ($p=0.013$) and BMI ($p=0.008$) compared with sedentary individuals. There was no significant difference in ABSI, Hcy and WC between healthy subjects with different activity levels.

Results of the t-test indicated a significant difference between sedentary CVD patients and sedentary healthy subjects for ABSI ($p=0.008$), BRI ($p=0.021$) and Hcy ($p=0.001$). Moreover, Hcy level differed significantly between moderately-active CVD patients and moderately-active healthy individuals ($p=0.039$). In addition, there were significant differences between active CVD patients and active healthy individuals in terms of BRI ($p=0.008$) and Hcy ($p=0.016$) levels.

Table 1- Mean values of research variables in middle-aged CVD patients and healthy individuals based on the level of physical

Variables	CVD patients				Healthy individuals			
	Sedentary	Moderately active	Active	P-value	Sedentary	Moderately active	Active	P-value
Age (years)	49.4 ± 4.7	44.2 ± 3.4	46.4 ± 5.4	0.047	51.2 ± 5.7	47.4 ± 4.4	49.2 ± 5.3	0.347
Weight (kg)	74.24±9.5	71.17±9.3	70.43±7.2	0.269	72.78±6.3	74.41±7.6	71.16±7.4	0.248
Height (cm)	162.2±5.1	161.67±6.2	164.14±5.1	0.147	163.1±4.5	165.3±7.4	162.87±6.2	0.311
ABSI	0.087±0.4	0.085±0.5	0.086±0.5	0.076	0.083±0.5	0.084±0.3	0.084±0.5	0.102
BRI	7.26 ± 1.7	6.85±1.4	6.08±2.1	0.031	6.14 ± 1.9	6.21±1.5	5.01±1.7	0.031
WC (cm)	99.39±8.5	96.36±9.7	95.14±8.3	0.147	100.2±9.3	95.11±8.7	94.25±7.9	0.034
BMI (kg/m ²)	28.42±3.8	27.68±4.1	26.25±3.1	0.089	27.53±4.2	27.34±3.8	26.85±5.3	0.167
Hcy (µmol/L)	18.87±2.1	19.25±1.7	16.26±2.3	0.008	14.51±2.4	13.14±2.9	13.42±2.1	0.089

Results are presented as mean ± standard deviation.

The basic characteristics and mean of variables of elderly subjects are shown in [table 2](#). Active CVD patients had significantly lower BRI ($p=0.041$), ABSI ($p=0.011$) and Hcy ($p=0.001$) levels compared with the sedentary counterparts. There was no significant difference in BMI and WC between the mentioned groups. Active healthy individuals had significantly lower BRI ($p=0.001$) and WC ($p=0.008$) levels compared with the sedentary counterparts. There was no significant difference in ABSI, WC and Hcy

between the mentioned groups.

Results of the t-test showed that there were significant differences between sedentary CVD patients and sedentary healthy subjects in terms of ABSI ($p=0.001$) and Hcy ($p=0.008$) levels. In addition, there were significant differences between moderately-active CVD patients and moderately-active healthy individuals in terms of Hcy ($p=0.001$) and WC ($p=0.049$). Moreover, WC differed significantly between active CVD patients and active healthy individuals ($p=0.001$).

Table 2- Mean values of research variables in old CVD patients and healthy individuals based on the level of physical activity

Variables	CVD patients			P-value	Healthy individuals			P-value
	Sedentary	Moderately active	Active		Sedentary	Moderately active	Active	
Age (years)	66.3 ± 5.7	68.1 ± 4.4	65.2 ± 5.1	0.0126	68.2 ± 6.1	64.1 ± 4.7	65.7 ± 5.6	0.214
Weight (kg)	72.31±6.4	72.1±7.2	71.48±7.1	0.347	74.5±7.1	71.82±5.2	73.23±6.1	0.187
Height (cm)	160.4±4.7	159.27±4.2	161.14±6.1	0.347	161.5±5.3	163.5±6.2	162.46±5.1	0.389
ABSI	0.089±0.4	0.087±0.5	0.086±0.5	0.008	0.086±0.5	0.085±0.5	0.085±0.4	0.214
BRI	7.84 ± 1.5	7.53±1.8	6.49±1.7	0.021	7.51 ± 1.7	7.11±1.6	5.77±1.6	0.011
WC (cm)	97.14±6.3	98.47±8.2	96.47±8.3	0.211	102.1±9.2	97.11±8.7	93.17±6.4	0.001
BMI (kg/m ²)	28.31±4.2	28.53±5.2	27.61±4.3	0.113	28.13±5.7	27.01±4.5	27.99±4.8	0.129
Hcy (µmol/L)	20.11±3.7	20.24±2.8	15.11±2.7	0.001	17.24±2.1	16.67±2.4	15.94±2.5	0.056

Results are presented as mean ± standard deviation

DISCUSSION

Based on the results, CVD patients had higher WC, BMI, Hcy, BRI and ABSI levels compared with healthy subjects. Both CVD patients and healthy individuals with a higher physical activity level had lower values of WC, BMI, Hcy, BRI and ABSI. These findings are consistent with results of a study by Vincent et al. that indicated a decrease in Hcy level following resistance training among inactive elderly people ([21](#)). Two other studies also showed that Hcy decreased after exercise training in male CVD patients and obese/overweight women ([22](#), [23](#)). On the other hand, Di Santolo et al. found no relationship between recreational sports and Hcy level among young women ([24](#)). Another study also showed that Hcy concentration is not significantly affected by submaximal exercise ([25](#)). Furthermore, Bambaiechi et al. reported that aerobic exercise did not change Hcy level in young ([26](#)). Various factors such as intensity, type and duration of exercise as well as gender and age of the participants may have contributed to the discrepancy between results of the mentioned studies.

Elevated level of Hcy has been associated with an increased risk of CVD. Plausible mechanisms of this association include vascular endothelial activation, promotion of oxidation of low-density lipoprotein cholesterol, vascular smooth cell proliferation, generation of superoxide anion and coagulation abnormalities ([27](#)). The effect of exercise on Hcy can be explored from different aspects. This amino acid is a critical biochemical juncture between methionine metabolism and the biosynthesis of the amino acids' cysteine and taurine; therefore, decreased methionine level leads to Hcy reduction ([27,28](#)). Increased demand for energy during exercise training might lead to the conversion of Hcy back to methionine, which can be used in the Krebs cycle for energy production. The relationship of Hcy with vitamin B12 has also been reported confirming the importance of these vitamins for regulating Hcy levels. In addition, reduction of oxidative stress indices following regular exercise can also reduce Hcy levels ([28](#)).

The present study showed that BRI and ABSI were significantly lower in physically-active subjects than in sedentary subjects. Some studies have found that BRI has a good discriminative power for diabetes (13) and CVD and its risk factors (29) compared to BMI, WC and other indices. Chang et al. showed that the adjusted odds ratio (OR) for predicting diabetes increased with increasing quintiles of BRI, after adjustment for age, smoking, alcohol consumption and other confounding variables (11). Despite the favorable discriminative capabilities of BRI, a study reported that the adjusted OR of BRI for health outcomes was not superior to that of BMI or WC in men and women, except for hyperuricemia in women (29). Consistent with results of some previous studies (29,30), the present study emphasized a coincident predictive ability between BRI and weight-to-height ratio with respect to area under the receiver operating characteristic and partial correlation coefficients for cardiometabolic abnormalities. The advantage of BRI over the weight-to-height ratio could be related to the ability of accurate estimation of body fat percentage and visceral adipose tissue (11). Several studies have delineated ABSI as an inferior indicator of the CVD risk when compared to indices such as waist-to-height ratio and WC. In line with these findings, an earlier study concluded that waist to-height ratio and WC are better predictors of CVD risk compared to ABSI, despite ABSI being directly proportional to WC (31). Savva et al. have specified that the risk factors of CVD in children can be better predicted by WC and waist-to-height ratio (32). Adegbija et al. indicated that the risk of CVD is directly proportional to WC (33). Similarly, Haghghatdoost et al. have concluded that ABSI is a weaker predictor of CVD risk in an Iranian population than BMI and waist to-height ratio (12). In contrast, Bozorgmanesh et al. has reported ABSI as a better indicator of CVD risk than BMI, WC and waist-to-hip ratio (34). Findings also suggest that ABSI is more significantly associated with circulating total cholesterol and insulin than BMI in young sedentary men. Regular physical activity is considered a cornerstone in the prevention and management of hypertension and CVD (35). It has been well-demonstrated that physical activity can reduce the risk of CVD-related morbidity and

mortality (36). However, the effect of physical activity might partly be mediated by adiposity (37), implying that the adjustment for WC is overly conservative. Physical activity of at least moderate intensity affects a range of biological processes that may influence cardiometabolic risk factors without affecting adiposity.

CONCLUSION

Our findings indicate that major CVD risk factors were favorably affected by physical activity. Active individuals generally had lower CVD risk factors. It is suggested to offer behavioral interventions to sedentary individual to promote the effectiveness of physical activity as a supportive mean for a healthier lifestyle.

ACKNOWLEDGMENTS

This article is derived from a PhD thesis approved by the Islamic Azad University, Aliabad Katool Branch, Iran. The authors would like to express their deep gratitude to all people who have collaborated in this study.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest regarding publication of this article.

REFERENCES

1. Roth GA, Johnson C, Abajobir A, Abd-Allah F, Abera SF, Abyu G, et al. *Global, regional, and national burden of cardiovascular diseases for 10 causes, 1990 to 2015. J Am Coll Cardiol.* 2017; 70(1): 1-25. [DOI:10.1016/j.jacc.2017.04.052] [PubMed] [Google Scholar]
2. World Health Organization. *Physical activity*[Online]. [cited 2012]; Available from: URL: http://www.who.int/topics/physical_activity/en. [Google Scholar]
3. Neeland IJ, Poirier P, Després J-P. *Cardiovascular and metabolic heterogeneity of obesity: clinical challenges and implications for management.* *Circulation.* 2018;137(13): 1391-406. [DOI:10.1161/CIRCULATIONAHA.117.029617] [PubMed] [Google Scholar]
4. Luo J, Hendryx M, Laddu D, Phillips LS, Chlebowski R, LeBlanc ES, et al. *Racial and ethnic differences in anthropometric measures as risk factors for diabetes.* *Diabetes Care.* 2019;42(1): 126-33. [DOI:10.2337/dc18-1413] [PubMed] [Google Scholar]
5. Berrington de Gonzalez A, Hartge P, Cerhan JR, Flint AJ, Hannan L, RJ MI, et al. *Body-mass index and mortality among 1.46 million white adults.* *New Engl J Med.* 2010;363(23): 221-9. [DOI:10.1056/NEJMoa1000367] [PubMed] [Google Scholar]

6. Zeng Q, He Y, Dong S, Zhao X, Chen Z, Song Z, et al. *Optimal cut-off values of BMI, waist circumference and waist: height ratio for defining obesity in Chinese adults.* *Br J Nutr.* 2014;112(10): 1735-44. [DOI:10.1017/S0007114514002657] [PubMed] [Google Scholar]
7. Donini LM, Poggiogalle E, Del Balzo V. *How to estimate fat mass in overweight and obese subjects.* *Int J Endocrinol.* 2013;28: 56-70. [DOI:10.1155/2013/285680] [PubMed] [Google Scholar]
8. Krakauer NY, Krakauer JC. *A new body shape index predicts mortality hazard independently of body mass index.* *PLoS One.* 2012; 7, 39-50. [DOI:10.1371/journal.pone.0039504] [PubMed] [Google Scholar]
9. He S, Chen X. *Could the new body shape index predict the new onset of diabetes mellitus in the Chinese population?* *PLoS One.* 2013; 8, 50-73. [DOI:10.1371/journal.pone.0050573] [PubMed] [Google Scholar]
10. Krakauer NY, Krakauer JC. *Dynamic association of mortality hazard with body shape.* *PLoS One.* 2014; 9, 88-93. [DOI:10.1371/journal.pone.0088793] [PubMed] [Google Scholar]
11. Chang Y, Guo X, Chen Y, Guo L, Li Z, Shasha Yu et al. *A body shape index and body roundness index: two new body indices to identify diabetes mellitus among rural populations in northeast China.* *BMC Public Health.* 2015; 15, 794-305. [DOI:10.1186/s12889-015-2150-2] [PubMed] [Google Scholar]
12. Haghghatdoost F, Sarrafzadegan N, Mohammadifard N, Asgary S, Boshtam M, Azadbakht L. *Assessing body shape index as a risk predictor for cardiovascular diseases and metabolic syndrome among Iranian adults.* *Nutrition.* 2014; 30, 636-644. [DOI:10.1016/j.nut.2013.10.021] [PubMed] [Google Scholar]
13. Thomas DM, Bredlau C, Bosy-Westphal A, Mueller M, Shen W, Gallagher D, et al. *Relationship between body roundness index with body fat and visceral adipose tissue emerging from a new geometrical model.* *Obesity (Silver Spring)* 2013; 21, 2264-2271. [DOI:10.1002/oby.20408] [PubMed] [Google Scholar]
14. Zhang N, Chang Y, Guo X, Chen Y, Ning Ye, Yingxian Sun. *A Body Shape Index and Body Roundness Index: two new body indices for detecting association between obesity and hyperuricemia in rural area of China.* *Eur J Intern Med.* 2016; 29, 32-36. [DOI:10.1016/j.ejim.2016.01.019] [PubMed] [Google Scholar]
15. McCully KS. *Homocysteine and the pathogenesis of atherosclerosis.* *Expert Review of Clinical Pharmacology,* 2015; 8(2):211-9. [DOI:10.1586/17512433.2015.1010516] [PubMed] [Google Scholar]
16. Chrysant SG, Chrysant GS. *The current status of homocysteine as a risk factor for cardiovascular disease: a mini review.* *Expert Review of Cardiovascular Therapy,* 2018, 16:8, 559-565. [DOI:10.1080/14779072.2018.1497974] [PubMed] [Google Scholar]
17. Dankner R, Geulayov G, Farber N, Novikov I, Segev S, Sela BA. *Cardiorespiratory fitness and plasma homocysteine levels in adult males and females.* *Isr Med Assoc J.* 2009 ;11(2):78-82 [PubMed] [Google Scholar]
18. Sinha D, Dwivedi MK. *The Effect of Physical Activity on Blood Homocysteine Concentration.* *Int J Curr Microbiol App Sci,* 2017, 6(9): 1206-10. [DOI:10.20546/ijcmas.2017.609.146] [Google Scholar]
19. Alomari MA, Khabour OF, Gharaibeh MY, Qhatan RA. *Effect of physical activity on levels of homocysteine, folate, and vitamin B12 in the elderly.* *Phys Sportsmed.* 2016;44(1):68-73. doi: 10.1080/00913847.2016.1135037. 2016. 1-6 [DOI:10.1080/00913847.2016.1135037] [PubMed] [Google Scholar]
20. Kelishadi R, Ardalan G, Gheiratmand R, Gouya MM, Razaghi EM, Delavari A, et al. *Association of physical activity and dietary behaviours in relation to the body mass index in a national sample of Iranian children and adolescents: CASPIAN Study.* *Bull World Health Organ.* 2007 ;85(1):19-26. [DOI:10.2471/BLT.06.030783] [PubMed] [Google Scholar]
21. Vincent KR, Braith RW, Bottiglieri T, Vincent HK, Lowenthal DT. *Homocysteine and lipoprotein levels following resistance training in older adults.* *Prev Cardiol.* 2003 Fall;6(4): 197-203. [DOI:10.1111/j.1520-037X.2003.01723.x] [PubMed] [Google Scholar]
22. Rostami S, Gaeini A, Zamani A, Hoseini H. *A Survey of the Effect of a Cardiac Rehabilitation Program (Concurrent Resistance-Endurance Training) on Homocysteine, Interleukin-6 and Interleukin-10 Changes in Men with Heart Disease.* *Sport physiology & management Investigations.* 2016, 8(2): 24-35. [Google Scholar]
23. Soori R, Choopani S, Falahian N. *Effect of Physical Activity on Serum Homocysteine Levels in Obese and Overweight Women.* *Quarterly of the Horizon of Medical Sciences,* 2017; 22(4): 307-312. [DOI:10.18869/acadpub.hms.22.4.307] [Google Scholar]
24. Di Santolo M, Banfi G, Stel G, Cauci S. *Association of recreational physical activity with homocysteine, folate and lipid markers in young women.* *European journal of applied physiology.* 2009; 105(1): 111-118. [DOI:10.1007/s00421-008-0880-x] [PubMed] [Google Scholar]
25. Subaşı SS, Gelecek N, Aksakoğlu G, Örmen M. *Effects of two different exercise trainings on plasma homocysteine levels and other cardiovascular disease risks.* *Turkish Journal of Biochemistry/Turk Biyokimya Dergisi.* 2012; 37(3), 61-74. [DOI:10.5505/tjb.2012.97268] [Google Scholar]
26. Bambaiechi E, Najari MA, Barjasteh B. *Influence of incremental aerobic exercise on homocysteine level in young males.* *British Journal of Sports Medicine.* 2010; 44(1): 21-29. [DOI:10.1136/bjism.2010.078725.70] [Google Scholar]
27. Hamedinia M, Haghghi A, Ravasi A. *The effect of aerobic training on inflammatory markers of cardiovascular disease risk in obese men.* *World J Sport Sci.* 2009; 2(1): 7-12. [Google Scholar]

28. Hmcic D, Rasic-Markovic A, Lekovic J, Krstic D, Colovic M, Macut D, et al. *Exercise decreases susceptibility to homocysteine seizures: the role of oxidative stress*. *Int J Sports Med*. 2014;35(7): 544-50. [DOI:10.1055/s-0033-1357162] [PubMed] [Google Scholar]
29. Chang Y, Guo X, Li T, Li S, Guo J, Sun Y. *A Body Shape Index and Body Roundness Index: Two New Body Indices to Identify left Ventricular Hypertrophy among Rural Populations in Northeast China*. *Heart Lung Circ*. 2016 ;25(4): 358-64 [DOI:10.1016/j.hlc.2015.08.009] [PubMed] [Google Scholar]
30. Maessen MF, Eijsvogels TM, Verheggen RJ, Hopman MT, Verbeek AL, de Vegt F. *Entering a new era of body indices: the feasibility of a body shape index and body roundness index to identify cardiovascular health status*. *PLoS One*. 2014 17;9(9):e107212. [DOI:10.1371/journal.pone.0107212] [PubMed] [Google Scholar]
31. Garcia-Hermoso A, Saavedra JM, Ramirez-Velez R, Ekelund U, Del Pozo-Cruz B. *Reallocating sedentary time to moderate-to-vigorous physical activity but not to light-intensity physical activity is effective to reduce adiposity among youths: a systematic review and meta-analysis*. *Obes Rev*. 2017; 18:1088-95. [DOI:10.1111/obr.12552] [PubMed] [Google Scholar]
32. Savva SC, Tornaritis M, Savva ME, Kourides Y, Panagi A, Silikiotou N, et al . *Waist circumference and waist-to-height ratio are better predictors of cardiovascular disease risk factors in children than body mass index*. *Int J Obes Relat Metab Disord*. 2000;24(11):1453-8. [DOI:10.1038/sj.ijo.0801401] [PubMed] [Google Scholar]
33. Adegbija O, Hoy W, Wang Z. *Prediction of cardiovascular disease risk using waist circumference among Aboriginals in a remote Australian community*. *BMC Public Health*. 2015; 15: 57-72. [DOI:10.1186/s12889-015-1406-1] [PubMed] [Google Scholar]
34. Bozorgmanesh M, Sardarinia M, Hajsheikholeslami F, Azizi F, Hadaegh F. *CVD-predictive performances of "a body shape index" versus simple anthropometric measures: Tehran lipid and glucose study*. *Eur J Nutr*. 2016; 55: 147-157. [DOI:10.1007/s00394-015-0833-1] [PubMed] [Google Scholar]
35. Skrede T, Stavnsbo M, Aadland E, Aadland KN, Anderssen SA, Resaland GK, et al. *Moderate-to-vigorous physical activity, but not sedentary time, predicts changes in cardiometabolic risk factors in 10-year-old children: the Active Smarter Kids Study*. *Am J Clin Nutr*. 2017 ;105(6):1391-1398. [DOI:10.3945/ajcn.116.150540] [PubMed] [Google Scholar]
36. Eckel RH, Jakicic JM, Ard JD, de Jesus JM, Houston Miller N, Hubbard VS, et al. *AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines*. *J Am Coll Cardiol*. 2014 1;63(25 Pt B):2960-84. [DOI:10.1161/01.cir.0000437740.48606.d1] [PubMed] [Google Scholar]
37. Pamidi S, Tasali E. *Obstructive sleep apnea and type 2 diabetes: is there a link?* *Front Neurol*. 2012; 3: 126-37. [DOI:10.3389/fneur.2012.00126] [PubMed] [Google Scholar]

How to Cite:

Arbabian T, Rezaeeshirazi R, Hesari A, Tavakol H [Does Physical Activity Level Affect Homocysteine and Obesity Variables in Women with Cardiovascular Disease?]. *mljgoums*. 2021; 15(4):21-27 DOI: 10.29252/mlj.15.4.21