

Interferon-Gamma Gene Polymorphism in Patients with Tuberculosis

Biranvand E (BSc)

Student of MSc, Immunology & microbiology
Department, Mazandaran University of Medical
Sciences

Abedian Kenary S (PhD)

Immunologist, Immunology & microbiology
Department, Molecular Biological Research
Center, Mazandaran University of Medical
Sciences

Ghaheeri A (MD)

General Practitioner of Health Center, Mazandaran
University of Medical Sciences

Rezaei M S (MD)

Infectious Disorders, Pediatrics Department of
Mazandaran University of Medical Sciences

Hasannia H(BSc)

Student of MSc, Immunology & microbiology
Department, Mazandaran University of Medical
Sciences

Nasrolahi M (PhD)

Professor of Microbiology, Faculty of Medical,
Mazandaran University of Medical Sciences

Parsaee MR (MD)

General Practitioner of Health Center, Mazandaran
University of Medical Sciences

Ahanjan M (PhD)

Microbiologist, Immunology & microbiology
Department, Mazandaran University of Medical
Sciences

Biranvand B (BSc)

Student of MSc of Biostatistics, Mazandaran
University of Medical Sciences

Ahmadi Basiri, E.

Medical Student, Ahvaz University of Medical
Sciences

Jivad F (BSc)

Student of MSc, Immunology & microbiology
Department, Mazandaran University of Medical
Sciences

Abstract

Background and objectives: Interferon-Gamma and interferon Gamma receptor (IFN γ / IFN γ R1) are the main genes associated with susceptibility to tuberculosis. We aimed at studying on interferon-Gamma Gene polymorphism(- 56 C/T) in people suffered from tuberculosis (TB).

Material and Methods: In this case-control study, the subjects were 62 individuals with TB and 74 healthy ones. Genomic DNA was extracted by DNA isolation kit(Roche Corporation), and genotype identification was performed by polymerase chain reaction-Restriction fragment length polymorphism (PCR-RFLP). Chi square and logistic regression, using SPSS soft ware (version 18), was used to compare genotype and alleles between case and control groups.

Results: The frequency of TT genotype in tuberculosis patients and healthy person are 43.5% and 17.5%, respectively. Based on Logistic regression (odd ration 0.148, p=0.0006), there is significant difference between Case and Control. In addition, the frequency of T allele is, in case group, 62.09 % the difference between case and control is significant, based on Logistic regression (odd ratio: 0.418, P=0.028).

Conclusion: It is implied that -56C/T is associated with IFN γ R1 promoter in tuberculosis patient. It is found to be associated with increased susceptibility to tuberculosis.

Key words: Tuberculosis, IFN γ R, PCR-RFLP

Corresponding Author: Abedian Kenary S
E-mail: abedianlab@yahoo.co.uk